arXiv:1209.1146v1 [math.AP] 5 Sep 2012

On linear instability of solitary waves for the nonlinear Dirac equation

Andrew Comech
Texas A&M University, College Station, TX 77843, U.S.A.
Institute for Information Transmission Problems, Moscow 101447, Russia

Meijiao Guan, Stephen Gustafson
University of British Columbia, Vancouver V6T 1Z2, Canada

September 2, 2012

Abstract. We consider the nonlinear Dirac equation, also known as the Soler model:

$$i\partial_t\psi = -i\boldsymbol{\alpha}\cdot\boldsymbol{\nabla}\psi + m\beta\psi - f(\psi^*\beta\psi)\beta\psi, \qquad \psi(x,t)\in\mathbb{C}^N, \qquad x\in\mathbb{R}^n, \qquad f\in C^2(\mathbb{R}),$$

where α_j, $j=1,\dots,n$, and β are $N\times N$ Hermitian matrices which satisfy $\alpha_j^2=\beta^2=I_N$, $\alpha_j\beta+\beta\alpha_j=0$, $\alpha_j\alpha_k+\alpha_k\alpha_j=2\delta_{jk}I_N$. We study the spectral stability of solitary wave solutions $\phi(x)e^{-i\omega t}$. More precisely, we study the presence of point eigenvalues in the spectra of linearizations at the small amplitude solitary waves in the limit $\omega\to m$. We consider dimensions $n=1$, 2, and 3, and prove that if $f(s)=s^k+O(s^{k+1})$, $k\in\mathbb{N}$, with $m>0$ and either $k\geq 3$, $n=1$, or $k\geq 2$, $n=2$, or $k=1$, $n=3$, then one positive and one negative eigenvalue are present in the spectrum of the linearizations at these solitary waves with ω sufficiently close to $\omega=m$. This shows in particular that these solitary waves are linearly unstable. The approach is based on applying the Rayleigh-Schrödinger perturbation theory to the nonrelativistic limit of the equation. The results are in formal agreement with the Vakhitov-Kolokolov stability criterion.

Résumé. Nous considérons l'équation de Dirac non linéaire, aussi connu comme le modèle de Soler:

$$i\partial_t\psi = -i\boldsymbol{\alpha}\cdot\boldsymbol{\nabla}\psi + m\beta\psi - f(\psi^*\beta\psi)\beta\psi, \qquad \psi(x,t)\in\mathbb{C}^N, \qquad x\in\mathbb{R}^n, \qquad f\in C^2(\mathbb{R}).$$

Nous étudions la stabilité spectrale des ondes solitaires solutions $\phi(x)e^{-i\omega t}$. Plus précisément, nous étudions la présence de valeurs propres dans les spectres des linéarisations aux ondes solitaires des petites amplitudes dans la limite $\omega\to m$. Nous considérons dimensions $n=1$, 2 et 3, et montrons que si $f(s)=s^k+O(s^{k+1})$, $k\in\mathbb{N}$, avec $m>0$ et soit $k\geq 3$, $n=1$, ou $k\geq 2$, $n=2$, ou $k=1$, $n=3$, ensuite une valeur propre positive et une négative sont présents dans le spectre des linéarisations à ces ondes solitaires lorsque ω est suffisamment proche de $\omega=m$. Ceci montre en particulier que ces ondes solitaires sont linéairement instable. L'approche est basée sur l'application de théorie de la perturbation de Rayleigh-Schrd̈inger à la limite non relativiste de l'équation. Les résultats sont en accord formel avec le critère de stabilité de Vakhitov-Kolokolov.

1 Introduction

A natural simplification of the Dirac-Maxwell system [Gro66] is the nonlinear Dirac equation, such as the massive Thirring model [Thi58] with vector-vector self-interaction and the Soler model [Sol70] with scalar-scalar self-interaction (known in dimension $n=1$ as the massive Gross-Neveu model [GN74, LG75]). There is an enormous body of research devoted to the nonlinear Dirac equation, which we can not cover comprehensively here. The existence of standing waves in the nonlinear Dirac equation was studied in [Sol70], [CV86], [Mer88], and [ES95]. Numerical confirmation of spectral stability of solitary waves of small amplitude is contained in [Chu07]. An overview of the well-posedness of the nonlinear Dirac equation in 1D is contained in [Pel10]. The asymptotic stability of small amplitude solitary waves in an external potential has been studied in [Bou08, PS10].

These models with self-interaction of local type have been receiving a lot of attention in particle physics, as well as in the theory of Bose-Einstein condensates [HC09, MJZ$^+$10]. The question of stability of solitary waves is of utmost importance: perturbations ensure that we only ever encounter stable configurations. Recent attempts at asymptotic stability [PS10, BC11] rely on the fundamental question of *spectral stability*:

> *Consider the Ansatz* $\psi(x,t) = \big(\phi_\omega(x) + \rho(x,t)\big)e^{-i\omega t}$, *with* $\phi_\omega(x)e^{-i\omega t}$ *a solitary wave solution. Let* $\partial_t\rho = A_\omega\rho$ *be the linearized equation for* ρ*. Does* A_ω *have eigenvalues in the right half-plane?*

In spite of a very clear picture of the spectral stability for nonlinear Schrödinger and Klein-Gordon equations [VK73, Sha83, Wei85, SS85, GSS87] and many attempts at spectral stability in the nonlinear Dirac context (let us mention [AS83, AS86, SV86, Bou08, CKMS10]), in the latter case the question of spectral stability is still completely open. Numerical results [BC09] show that in the 1D Soler model (cubic nonlinearity) all solitary waves are spectrally stable. Let us mention the related results [CP06, Chu07].

Previous results [Com11] show that the Vakhitov-Kolokolov criterion in the case of the nonlinear Dirac equation gives a less definite answer about the spectral stability than in the case of the nonlinear Schrödinger equation. All we know is that at the value of ω where $\partial_\omega Q(\omega)$ vanishes, with $Q(\omega)$ being the charge of the solitary wave $\phi_\omega e^{-i\omega t}$, two eigenvalues of the linearized equation collide at $\lambda = 0$, but we do not know where these eigenvalues are located when $\partial_\omega Q(\omega) \neq 0$. This is in contrast to the ground states of the nonlinear Schrödinger equation, when the condition $\partial_\omega Q(\omega) > 0$ is enough to conclude that one positive and one negative eigenvalue of the linearized operator move out of $\lambda = 0$ along the real axis; see [VK73, GSS87, CP03].

In the present paper, we show that if Vakhitov-Kolokolov guarantees linear instability for the system obtained in the nonrelativistic limit, then the same result is true for the corresponding small amplitude solitary waves in the original relativistic system. We use our approach to show that the small amplitude solitary wave solutions of the nonlinear Dirac equation are linearly unstable if the Vakhitov-Kolokolov condition says so. Let us mention that our results apply to the solitary wave solutions which we obtain from the solitary waves of the nonlinear Schrödinger equation in the nonrelativistic limit of the nonlinear Dirac.

According to our results, the spectrum of the linearization at small amplitude solitary waves in the 1D nonlinear Dirac equation with cubic and quintic nonlinearities has no real eigenvalues; instead, one can prove the existence of two purely imaginary eigenvalues. To prove that these solitary waves are indeed spectrally stable, one also needs to prove that there are no complex eigenvalues with $\operatorname{Re}\lambda > 0$; we already have partial results which we will publish elsewhere.

Our approach is based on the idea that the family of real eigenvalues of the linearization at a solitary wave of the nonlinear Dirac equation bifurcating from $\lambda = 0$ is a deformed family of eigenvalues of the corresponding linearization of the nonlinear Schrödinger equation. The model and the main results are described in Section 2. The necessary constructions in the context of the nonlinear Schrödinger equation are presented in Section 3. The existence, asymptotics, and linearization of solitary waves of the nonlinear Dirac equation is covered in Section 4. The statement of Theorem 2.1 for $\omega \lesssim m$ follows from Proposition 5.2, which we prove using the Rayleigh-Schrödinger perturbation theory.

2 Main result

We consider the nonlinear Dirac equation

$$i\partial_t\psi = -i\boldsymbol{\alpha}\cdot\boldsymbol{\nabla}\psi + m\beta\psi - f(\psi^*\beta\psi)\beta\psi, \qquad \psi(x,t)\in\mathbb{C}^N, \qquad x\in\mathbb{R}^n, \tag{2.1}$$

with $m > 0$ and ψ^* being the Hermitian conjugate of ψ. We assume that the nonlinearity $f(s)$ is smooth and real-valued, and that

$$f(0) = 0.$$

Above, $\boldsymbol{\alpha}\cdot\boldsymbol{\nabla} = \sum\limits_{j=1}^{n} \alpha_j \frac{\partial}{\partial x_j}$, and the Hermitian matrices α_j and β are chosen so that

$$\big(-i\boldsymbol{\alpha}\cdot\boldsymbol{\nabla} + \beta m\big)^2 = (-\Delta + m^2) I_N,$$

where I_N is the $N\times N$ unit matrix. That is, α_j and β are to satisfy

$$\alpha_j\alpha_k + \alpha_k\alpha_j = 2\delta_{jk} I_N, \qquad \beta^2 = I_N; \qquad \alpha_j\beta + \beta\alpha_j = 0. \tag{2.2}$$

The generalized massive Gross-Neveu model, or, in the terminology of [CKMS10], the scalar-scalar case with $k\in\mathbb{N}$, corresponds to the nonlinearity $f(s) = s^k$.

According to the Dirac-Pauli theorem (cf. [Dir28, vdW32, Pau36] and [Tha92, Lemma 2.25]), the particular choice of the matrices α_j and β does not matter. In terms of the standard Pauli matrices,

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \qquad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \qquad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \tag{2.3}$$

which satisfy $\sigma_i\sigma_j + \sigma_j\sigma_i = 2\delta_{ij} I_2$, we take:

$$n=1: \quad N=2,\ \alpha = -\sigma_2,$$

$$n=2: \quad N=2,\ \alpha_j = \sigma_j, \quad 1\le j\le 2,$$

$$n=3: \quad N=4,\ \alpha_j = \begin{pmatrix} 0 & \sigma_j \\ \sigma_j & 0 \end{pmatrix}, \quad 1\le j\le 3,$$

and in all cases

$$\beta = \begin{pmatrix} I_{N/2} & 0 \\ 0 & -I_{N/2} \end{pmatrix}.$$

For a large class of nonlinearities $f(s)$, there are solitary wave solutions of the form

$$\psi(x,t) = \phi_\omega(x) e^{-i\omega t}, \qquad \phi_\omega \in H^1(\mathbb{R}, \mathbb{C}^N), \qquad |\omega| < m, \tag{2.4}$$

where in dimensions $n = 1$, 2, and 3 respectively,

$$\phi_\omega(x) = \begin{bmatrix} v(x,\omega) \\ u(x,\omega) \end{bmatrix}, \quad \phi_\omega(x) = \begin{bmatrix} v(r,\omega) \\ i e^{i\phi} u(r,\omega) \end{bmatrix}, \quad \phi_\omega(x) = \begin{bmatrix} v(r,\omega)\begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ iu(r,\omega)\begin{pmatrix} \cos\theta \\ e^{i\phi}\sin\theta \end{pmatrix} \end{bmatrix}. \tag{2.5}$$

For $n = 1$, $v(x,\omega)$ can be taken positive and even, and $u(x,\omega)$ real-valued and odd; under these conditions, the solitary wave $\phi_\omega(x)$ is unique (see Section 4.1 for details).

For $n = 2$ and $n = 3$, $v(r,\omega)$ and $u(r,\omega)$ are real-valued, radially-symmetric functions, (r,ϕ) are standard polar coordinates in $\mathbb{R}^2$, and (r,θ,ϕ) are standard spherical coordinates in $\mathbb{R}^3$. The existence of solitary waves of this form is proved, for example, in [CV86]. The particular solutions we consider here, however, are those constructed for ω close to m, as outlined in Section 4.2, from the nonrelativistic limit $\omega\to m$.

Due to the $\mathbf{U}(1)$-invariance, for solutions to (2.1) the value of the charge functional

$$Q(\psi) = \int_{\mathbb{R}^n} |\psi(x,t)|^2\,dx$$

is formally conserved. For brevity, we also denote by $Q(\omega)$ the charge of the solitary wave $\phi_\omega(x)e^{-i\omega t}$:

$$Q(\omega) = \int_{\mathbb{R}^n} |\phi_\omega(x)|^2\,dx. \tag{2.6}$$

We are interested in the spectral stability of linearization of (2.1) at a solitary wave solution (2.4).

Theorem 2.1. *Let $n \le 3$. Assume that $f(s) = s^k$, where $k \in \mathbb{N}$ satisfies $k \ge 3$ for $n = 1$, or $k \ge 2$ for $n = 2$, or $k = 1$ for $n = 3$. Then the solitary wave solutions $\phi_\omega e^{-i\omega t}$ to (2.1) described above are linearly unstable for $\omega \in I$, where I is the largest interval with $\sup I = m$ such that $\partial_\omega Q(\omega)$ does not vanish on I. More precisely, let A_ω be the linearization of the nonlinear Dirac equation at a solitary wave $\phi_\omega(x)e^{-i\omega t}$. Then there are eigenvalues*

$$\pm\lambda_\omega \in \sigma_p(A_\omega), \qquad \lambda_\omega > 0 \quad \text{for } \omega \in I, \qquad \lambda_\omega = O(m-\omega).$$

See Figure 1.



Figure 1: Main result: The point spectrum of the linearization of the nonlinear Dirac equation in $\mathbb{R}^n$, $n \le 3$, with $f(s) = s^k$, $k \ge 4-n$, at a solitary wave with $\omega \lesssim m$ contains two nonzero real eigenvalues, $\pm\lambda_\omega$, with $\lambda_\omega = O(m-\omega)$. See Theorem 2.1. Also plotted on this picture is the essential spectrum, with the edges at $\lambda = \pm i(m-\omega)$ and with the embedded threshold points (branch points of the dispersion relation) at $\lambda = \pm i(m+\omega)$.

Remark 2.2. Theorem 2.1 extends easily to nonlinearities $f \in C^2(\mathbb{R})$ of the form

$$f(s) = as^k + O(s^{k+1}), \qquad a > 0.$$

Remark 2.3. We only need to prove the linear instability for $\omega \lesssim m$. Then, by [Com11], the positive and negative eigenvalues remain trapped on the real axis, not being able to collide at $\lambda = 0$ for $\omega \in I = (\omega_0, m)$ as long as $\partial_\omega Q(\omega)$ does not vanish on I (this nonvanishing is a sufficient condition for the dimension of the generalized null space to remain equal to two). These eigenvalues cannot leave into the complex plane, either, since they are simple, while the spectrum of the operator is symmetric with respect to the real and imaginary axes.

Remark 2.4. We do not know what happens at $\omega = \inf I$: as ω drops below $\inf I$, it could be that either the pair of real eigenvalues, having collided at $\lambda = 0$ when $\omega = \inf I$, turn into a pair of purely imaginary eigenvalues (linear instability disappears), or instead two purely imaginary eigenvalues, having met at $\lambda = 0$, turn into the second pair of real eigenvalues (linear instability persists).

Remark 2.5. Theorem 2.1 is in formal agreement with the Vakhitov-Kolokolov stability criterion [VK73], since for $\omega \lesssim m$ one has $Q'(\omega) > 0$ for $k \ge 4-n$. Let us mention that the sign of the instability criterion, $Q'(\omega) > 0$ differs from [VK73] because of their writing the solitary waves in the form $\varphi(x)e^{+i\omega t}$.

Remark 2.6. We expect that in the 1D case with $k = 1$ the small amplitude solitary waves are spectrally stable; we will prove this elsewhere.

We also expect that in the 1D case with $k = 2$ ("quintic nonlinearity", $f(\psi^*\beta\psi)\beta\psi = O(|\psi|^5)$) the small solitary wave solutions of the nonlinear Dirac equation in 1D are spectrally stable. For the corresponding nonlinear Schrödinger equation (quintic nonlinearity in 1D), the charge is constant, thus the zero eigenvalue of a linearized operator is always of higher algebraic multiplicity. For the Dirac equation, this degeneracy is "resolved": the charge is now a decaying function for $\omega \lesssim m$ (with nonzero limit as $\omega \to m$), suggesting that there are two purely imaginary eigenvalues $\pm\lambda_\omega$ in the spectrum of A_ω, with $\lambda_\omega = o(m-\omega)$, but no eigenvalues with nonzero real part.

Remark 2.7. Let us notice that in the 3D case for the cubic nonlinearity $f(s) = s$ (this is the original Soler model from [Sol70]), based on the numerical evidence from [Sol70, AS83], one expects that the charge $Q(\omega)$ has a local minimum at $\omega = 0.936$, suggesting that the solitary waves with $0.936 < \omega < 1$ are linearly unstable, but then at $\omega = 0.936$ the real eigenvalues collide at $\lambda = 0$, and there are no nonzero real eigenvalues in the spectrum for $\omega \lesssim 0.936$.

Remark 2.8. We can not rule out the possibility that the eigenvalues with nonzero real part could bifurcate directly from the imaginary axis into the complex plane. Such a mechanism is absent for the nonlinear Schrödinger equation linearized at a solitary wave, for which the point eigenvalues always remain on the real or imaginary axes. At present, though, we do not have examples of such bifurcations in the context of nonlinear Dirac equation linearized at a solitary wave.

Remark 2.9. For $n = 3$, we only consider the case $k = 1$, and we do not consider dimensions $n > 3$. This is because of the fact that the equation $-\Delta u + u = |u|^{p-1}u$ in $\mathbb{R}^n$ has nontrivial solutions in $H^1(\mathbb{R}^n)$ if and only if $1 < p < \frac{n+2}{n-2}$ [BL83, Example 1]. This does not allow us to construct solitary wave solutions to the nonlinear Dirac equation in $\mathbb{R}^3$ with $k > 2$ and in $\mathbb{R}^n$, $n > 3$, with any $k \in \mathbb{N}$.

3 Nonlinear Schrödinger and its solitary waves

We are going to use the fact that the nonlinear Dirac equation in the nonrelativistic limit coincides with the nonlinear Schrödinger equation,

$$i\partial_t\psi = -\frac{1}{2m}\Delta\psi - |\psi|^{2k}\psi \qquad \psi(x,t) \in \mathbb{C}, \quad x \in \mathbb{R}^n, \quad n = 1,\, 2,\, 3. \tag{3.1}$$

3.1 Solitary waves

The properties of solitary wave solutions

$$\psi(x,t) = \phi_\omega(x)e^{-i\omega t}, \qquad \phi_\omega \in H^1(\mathbb{R}^n),$$

with amplitude $\phi_\omega(x)$ therefore satisfying the equation

$$-\frac{1}{2m}\Delta\phi(x) - |\phi|^{2k}\phi = \omega\phi, \qquad x \in \mathbb{R}^n, \tag{3.2}$$

are well-known [Str77, BL83]. For any $k \in \mathbb{N}$ when $n \le 2$ and for $k = 1$ when $n = 3$, and for any $\omega \in (-\infty, 0)$, there is a unique positive, radially symmetric solution

$$\phi_\omega(x) = \phi_\omega(|x|) > 0,$$

which decays exponentially. This family of solitary waves, known as the *ground states*, is generated by scalings of a single amplitude function:

$$\phi_\omega(x) = |\omega|^{\frac{1}{2k}} F(\sqrt{2m|\omega|}x), \qquad \omega < 0, \tag{3.3}$$

where $F(x) = F(|x|) > 0$ solves

$$-\Delta F - F^{2k+1} = -F, \qquad x \in \mathbb{R}^n. \tag{3.4}$$

In one space dimension ($n = 1$), for $k \in \mathbb{N}$, there is the explicit formula

$$F(x) = \left(\frac{k+1}{\cosh^2 kx}\right)^{\frac{1}{2k}}.$$

3.2 Linearization at a solitary wave

To derive the linearization of the nonlinear Schrödinger equation (3.1) at a solitary wave $\psi(x,t)=\phi_\omega(x)e^{-i\omega t}$, we use the Ansatz

$$\psi(x,t)=(\phi_\omega(x)+\rho(x,t))e^{-i\omega t},\qquad \rho(x,t)\in\mathbb{C},\qquad x\in\mathbb{R}^n,$$

and arrive at the linearized equation

$$\partial_t\boldsymbol{\rho}=\mathbf{j}\mathbf{l}(\omega)\boldsymbol{\rho},\qquad \boldsymbol{\rho}(x,t)=\begin{bmatrix}\operatorname{Re}\rho(x,t)\\ \operatorname{Im}\rho(x,t)\end{bmatrix}, \tag{3.5}$$

where

$$\mathbf{j}=\begin{bmatrix}0&1\\-1&0\end{bmatrix},\qquad \mathbf{l}=\begin{bmatrix}\mathfrak{l}_-&0\\0&\mathfrak{l}_+\end{bmatrix}, \tag{3.6}$$

with $\mathfrak{l}_\pm$ self-adjoint Schrödinger operators

$$\mathfrak{l}_-(\omega)=-\frac{1}{2m}\Delta+|\phi_\omega|^{2k}-\omega,\qquad \mathfrak{l}_+(\omega)=\mathfrak{l}_-(\omega)-2k|\phi_\omega|^{2k}. \tag{3.7}$$

Since the solitary wave amplitudes $\phi_\omega(x)=\phi_\omega(|x|)$ we take here are radially symmetric, we may consider the operators

$$\mathbf{l}_{rad},\ \mathfrak{l}_{\pm,rad}\ :=\ \mathbf{l},\ \mathfrak{l}_\pm \text{ restricted to radially symmetric functions.} \tag{3.8}$$

The linear stability theory of NLS ground states is well understood, and can be summarized, in terms of their charge (2.6) as follows:

Lemma 3.1 (Vakhitov-Kolokolov stability criterion [VK73]). *For the linearization* (3.5) *at a ground state solitary wave* $\phi_\omega(x)e^{-i\omega t}$, *there are real nonzero eigenvalues* $\pm\lambda\in\sigma_d(\mathbf{j}\mathbf{l})$, $\lambda>0$, *if and only if* $\frac{d}{d\omega}Q(\omega)>0$ *at this value of* ω. *If so, then* $\pm\lambda\in\sigma_d(\mathbf{j}\mathbf{l}_{rad})$ *are simple eigenvalues, and moreover* $\ker\mathfrak{l}_{+,rad}=\{0\}$.

Using (3.3),

$$Q(\omega)=\int_{\mathbb{R}^n}|\phi_\omega(x)|^2\,dx=|\omega|^{1/k}\int_{\mathbb{R}^n}F^2(\sqrt{2m|\omega|}x)\,dx=C|\omega|^{1/k-n/2},\qquad \omega<0, \tag{3.9}$$

where $C=\int_{\mathbb{R}^n}F^2(\sqrt{2m}y)\,dy>0$. We see from (3.9) that for $\omega<0$, $Q'(\omega)<0$ for $k<2/n$, $Q'(\omega)=0$ for $k=2/n$, and $Q'(\omega)>0$ for $k>2/n$. Thus:

Lemma 3.2. *Let* $n\le 3$. *If* $k\in\mathbb{N}$ *satisfies* $k\ge 4-n$, *then* $\sigma_p(\mathbf{j}\mathbf{l}_{rad})\ni\{\pm\lambda\}$, *for some* $\lambda>0$, *and in particular the NLS ground states are linearly unstable.*

4 Nonlinear Dirac and its solitary waves

Solitary waves are solutions to (2.1) of the form

$$\psi(x,t)=\phi_\omega(x)e^{-i\omega t};\ \phi_\omega\in H^1(\mathbb{R},\mathbb{C}^N),\ \omega\in\mathbb{R}$$

and as such the amplitude $\phi_\omega(x)$ must satisfy

$$\omega\phi_\omega=-i\boldsymbol{\alpha}\cdot\boldsymbol{\nabla}\psi+m\beta\psi-f(\psi^*\beta\psi)\beta\psi,\qquad x\in\mathbb{R}^n. \tag{4.1}$$

4.1 Solitary waves in one dimension

We first give a simple demonstration of the existence and uniqueness of solitary waves in one dimension, following the article [BC09], and allowing for more general nonlinearities $f(s)$.

Lemma 4.1. *Let* $f(0) = 0$. *Denote* $g(s) = m - f(s)$, *and let* $G(s)$ *be the antiderivative of* $g(s)$ *such that* $G(0) = 0$. *Assume that there is* $\omega_0 < m$ *such that for given* $\omega \in (\omega_0, m)$ *there exists* $\varGamma_\omega > 0$ *such that*

$$\omega \varGamma_\omega = G(\varGamma_\omega), \qquad \omega \ne g(\varGamma_\omega), \qquad \text{and} \qquad \omega s < G(s) \quad \text{for } s \in (0, \varGamma_\omega). \tag{4.2}$$

Then there is a solitary wave solution $\psi(x,t) = \phi_\omega(x) e^{-i\omega t}$ *to* (2.1), *where*

$$\phi_\omega(x) = \begin{bmatrix} v(x,\omega) \\ u(x,\omega) \end{bmatrix}, \tag{4.3}$$

with both v *and* u *real-valued, belonging to* $H^1(\mathbb{R})$ *as functions of* x, v *being even and* u *odd.*

More precisely, for $x \in \mathbb{R}$ *and* $\omega \in (\omega_0, m)$, *let us define* $\mathscr{X}(x,\omega)$ *and* $\mathscr{Y}(x,\omega)$ *by*

$$\mathscr{X} = v^2 - u^2, \qquad \mathscr{Y} = vu. \tag{4.4}$$

Then $\mathscr{X}(x,\omega)$ *is the unique positive symmetric solution to*

$$\partial_x^2 \mathscr{X} = -\partial_{\mathscr{X}}(-2G(\mathscr{X})^2 + 2\omega^2 \mathscr{X}^2), \qquad \lim_{x\to\pm\infty} \mathscr{X}(x,\omega) = 0, \tag{4.5}$$

and $\mathscr{Y}(x,\omega) = -\frac{1}{4\omega}\partial_x \mathscr{X}(x,\omega)$. *This solution satisfies* $\mathscr{X}(0,\omega) = \varGamma_\omega$.

Proof. From (4.1), we obtain:

$$\begin{cases} \omega v = \partial_x u + g(|v|^2 - |u|^2) v, \\ \omega u = -\partial_x v - g(|v|^2 - |u|^2) u. \end{cases} \tag{4.6}$$

Assuming that both v and u are real-valued (this will be justified once we have found real-valued v and u), we can rewrite (4.6) as the following Hamiltonian system:

$$\begin{cases} \partial_x u = \omega v - g(v^2 - u^2) v = \partial_v h(v,u), \\ -\partial_x v = \omega u + g(v^2 - u^2) u = \partial_u h(v,u), \end{cases} \tag{4.7}$$

where the Hamiltonian $h(v,u)$ is given by

$$h(v,u) = \frac{\omega}{2}(v^2 + u^2) - \frac{1}{2} G(v^2 - u^2). \tag{4.8}$$

The solitary wave with a particular $\omega \in (\omega_0, m)$ corresponds to a trajectory of this Hamiltonian system such that

$$\lim_{x\to\pm\infty} v(x,\omega) = \lim_{x\to\pm\infty} u(x,\omega) = 0,$$

hence $\lim_{x\to\pm\infty} \mathscr{X} = 0$. Since $G(s)$ satisfies $G(0) = 0$, we conclude that $h(v(x), u(x)) \equiv 0$, which leads to

$$\omega(v^2 + u^2) = G(v^2 - u^2). \tag{4.9}$$

We conclude from (4.9) that solitary waves may only correspond to $|\omega| < m$, $\omega \ne 0$.

The functions $\mathscr{X}(x,\omega)$ and $\mathscr{Y}(x,\omega)$ introduced in (4.4) are to solve

$$\begin{cases} \partial_x \mathscr{X} = -4\omega \mathscr{Y}, \\ \partial_x \mathscr{Y} = -(v^2 + u^2) g(\mathscr{X}) + \omega \mathscr{X} = -\frac{1}{\omega} G(\mathscr{X}) g(\mathscr{X}) + \omega \mathscr{X}, \end{cases} \tag{4.10}$$

and to have the asymptotic behavior $\lim_{|x|\to\infty} \mathscr{X}(x) = 0$, $\lim_{|x|\to\infty} \mathscr{Y}(x) = 0$. In the second equation in (4.10), we used the relation (4.9). The system (4.10) can be written as the following equation on $\mathscr{X}$:

$$\partial_x^2 \mathscr{X} = -\partial_{\mathscr{X}}(-2G(\mathscr{X})^2 + 2\omega^2 \mathscr{X}^2) = 4\big(G(\mathscr{X}) g(\mathscr{X}) - \omega^2 \mathscr{X}\big). \tag{4.11}$$

This equation describes a particle in the potential $-2G(s)^2 + 2\omega^2 s^2$. The condition (4.2) is needed for the existence of the turning point of the zero energy trajectory in this potential at the point $s = \varGamma_\omega$. □

4.2 Solitary waves in the nonrelativistic limit

In dimensions $n = 1$, 2 and 3, we consider solitary wave amplitudes $\phi_\omega(x)$ of the forms given in (2.5). Substituting these into the nonlinear Dirac equation (2.1), a straightforward calculation results in the system

$$\left\{ \begin{array}{l} \omega v = \partial_r u + \frac{n-1}{r} u + mv - f(v^2 - u^2)v \\ \omega u = -\partial_r v - mu + f(v^2 - u^2)u \end{array} \right. \tag{4.12}$$

for the pair of real-valued functions $v = v(r, \omega)$, $u = u(r, \omega)$. Notice that equation (4.12) includes the $n = 1$-dimensional case (4.6) if we interpret $r = x$.

Recalling that $f(s) = s^k$, we arrive at

$$(\omega - m)v = \partial_r u + \frac{n-1}{r} u - fv, \qquad (\omega + m)u = -\partial_r v + fu,$$

with

$$f := (v^2 - u^2)^k.$$

For the nonrelativistic limit, we set

$$m^2 - \omega^2 = \epsilon^2, \quad 0 < \epsilon \ll m,$$

and rescale $v(r, \omega)$ and $u(r, \omega)$ as

$$v(r, \omega) = \epsilon^{\frac{1}{k}} V(\epsilon r, \epsilon), \qquad u(r, \omega) = \epsilon^{1+\frac{1}{k}} U(\epsilon r, \epsilon).$$

Then V, U should satisfy

$$\left\{ \begin{array}{l} (\omega - m)\epsilon^{\frac{1}{k}} V = \epsilon^{2+\frac{1}{k}} (\partial_R U + \frac{n-1}{R} U) - \epsilon^{1/k} fV \\ (\omega + m)\epsilon^{1+\frac{1}{k}} U = -\epsilon^{1+\frac{1}{k}} \partial_R V + \epsilon^{1+\frac{1}{k}} fU \end{array} \right. ,$$

where $R = \epsilon r$ denotes the "rescaled variable". Using $\omega = m - \frac{1}{2m}\epsilon^2 + O(\epsilon^4)$, and taking into account that

$$f = (\epsilon^{\frac{2}{k}} V^2 - \epsilon^{2+\frac{2}{k}} U^2)^k = \epsilon^2 V^{2k} + \epsilon^4 O(U^{2k} + V^{2k}),$$

we re-write the system as

$$\left\{ \begin{array}{l} (-\frac{1}{2m} + O(\epsilon^2))V = \partial_R U + \frac{n-1}{R} U - V^{2k+1} + \epsilon^2 O((U^{2k} + V^{2k})|V|) \\ (2m + O(\epsilon^2))U = -\partial_R V + \epsilon^2 V^{2k} U + \epsilon^4 O((U^{2k} + V^{2k})|U|) \end{array} \right. . \tag{4.13}$$

The rescaled system (4.13) has an obvious limit. Formally (for now) setting $\hat{V}(r) = \lim_{\epsilon\to 0} V(r, \epsilon)$, $\hat{U}(r) = \lim_{\epsilon\to 0} U(r, \epsilon)$, we arrive at

$$-\frac{1}{2m}\hat{V} = \partial_R \hat{U} + \frac{n-1}{R}\hat{U} - \hat{V}^{2k+1}, \qquad 2m\hat{U} = -\partial_R \hat{V}. \tag{4.14}$$

Substituting the second equation into the first yields

$$-\frac{1}{2m}(\partial_R^2 + \frac{n-1}{R}\partial_R)\hat{V} - \hat{V}^{2k+1} = -\frac{1}{2m}\hat{V}, \qquad \hat{U} = -\frac{1}{2m}\partial_R \hat{V}.$$

This equation for $\hat{V}(r)$ is precisely the equation (3.2) for NLS solitary wave amplitudes ϕ_ω with $\omega = -\frac{1}{2m}$. Thus we let $\hat{V}(r)$ be the (unique) NLS ground state:

$$\hat{V}(r) := (2m)^{-\frac{1}{2k}} F(r), \qquad \hat{U}(r) := -(2m)^{-\frac{1}{2k}-1} F'(r), \tag{4.15}$$

with $F(r)$ the unique positive spherically symmetric solution to (3.4). We can use this nonrelativistic limit to construct nonlinear Dirac solitary waves for $\epsilon^2 = m^2 - \omega^2 \ll m^2$:

Lemma 4.2. *There is* $\omega_0 < m$ *such that for* $\omega = \sqrt{m^2 - \epsilon^2} \in (\omega_0, m)$*, there are solutions of* (4.12) *of the form*

$$v(r,\omega) = \epsilon^{\frac{1}{k}} \left[\hat{V}(\epsilon r) + \tilde{V}(\epsilon r) \right], \quad u(r,\omega) = \epsilon^{1+\frac{1}{k}} \left[\hat{U}(\epsilon r) + \tilde{U}(\epsilon r) \right], \quad \|\tilde{V}\|_{H^2} + \|\tilde{U}\|_{H^2} = O(\epsilon^2).$$

Remark 4.3. In the one-dimensional case $n = 1$, since the solitary waves are unique (up to symmetries), it follows that these asymptotics describe *every* solitary wave for ω close to 1, or equivalently every small amplitude solitary wave.

Proof. The argument parallels that of [Gua08], where the (more general) nonlinearity $f(s) = |s|^{2\theta}$, $0 < \theta < 2$ is considered for $n = 3$. Writing

$$V(R,\epsilon) = \hat{V}(R) + \tilde{V}(R,\epsilon), \qquad V(R,\epsilon) = \hat{V}(R) + \tilde{V}(R,\epsilon),$$

and subtracting equations (4.13) and (4.14), we arrive at

$$\begin{aligned}
&-\frac{1}{2m}\tilde{V} + O(\epsilon^2)V \\
&\qquad = (\partial_R + \frac{n-1}{R})\tilde{U} - (2k+1)\hat{V}^{2k}\tilde{V} + O(|\hat{V}|^{2k+1}\tilde{V}^2 + |\tilde{V}|^{2k+1}) + \epsilon^2 O((U^{2k} + V^{2k})|V|), \\
&2m\tilde{U} + O(\epsilon^2)U = -\partial_R\tilde{V} + \epsilon^2 V^{2k}U + \epsilon^4 O((U^{2k} + V^{2k})|U|),
\end{aligned}$$

which, setting

$$\Xi(R,\epsilon) := \begin{bmatrix} \tilde{V}(R,\epsilon) \\ \tilde{U}(R,\epsilon) \end{bmatrix},$$

we may re-write as the system

$$\mathbf{H}\Xi = O_{H^1}(\epsilon^2) + O(\epsilon^2|\Xi| + |\Xi|^2 + |\Xi|^{2k+1}),$$

where

$$\mathbf{H} := \begin{bmatrix} -\frac{1}{2m} + (2k+1)\hat{V}^{2k} & -(\partial_R + \frac{n-1}{R}) \\ \partial_R & 2m \end{bmatrix}.$$

Since

$$\begin{bmatrix} \xi \\ \eta \end{bmatrix} \in \ker \mathbf{H} \iff \xi \in \ker \mathfrak{l}_+, \;\; \eta = -\frac{1}{2m}\partial_R \xi,$$

and $\ker \mathfrak{l}_{+,rad} = \{0\}$ (Cf. definitions (3.7), (3.8)), we see that $\ker \mathbf{H} = \{0\}$. It then follows from the fact that $\mathfrak{l}_{+,rad}^{-1}$ is bounded from $L^2_r(\mathbb{R}^n, \mathbb{C})$ to $H^2(\mathbb{R}^n, \mathbb{C})$, that $\mathbf{H}^{-1}$ is bounded from $H^1_r(\mathbb{R}^n, \mathbb{C}^2)$ to $H^2(\mathbb{R}^n, \mathbb{C}^2)$ (here L^2_r, H^1_r are the corresponding subspaces of spherically symmetric functions). Hence

$$\Xi = \mathbf{H}^{-1}\left\{ O_{H^1}(\epsilon^2) + O(\epsilon^2|\Xi| + |\Xi|^2 + |\Xi|^{2k+1}) \right\}, \tag{4.16}$$

and since $\|\Xi\|_{L^\infty} \leq C\|\Xi\|_{H^2}$ (recall $n \leq 3$), we arrive easily at

$$\begin{aligned}
\|R.H.S.(4.16)\|_{H^2} &\leq C\|O_{H^1}(\epsilon^2) + O(\epsilon^2|\Xi| + |\Xi|^2 + |\Xi|^{2k+1})\|_{H^1} \\
&\leq C\left\{ \epsilon^2 + \epsilon\|\Xi\|_{H^2} + \|\Xi\|^2_{H^2} + \|\Xi\|^{2k+1}_{H^2} \right\},
\end{aligned}$$

and so we see that for small enough ϵ, the map on the r.h.s. of (4.16) maps the ball of radius ϵ in H^2 into itself. A similar estimate shows that this map is a contraction, and hence has a unique fixed point Ξ in this ball. Finally, we see from (4.16) that $\|\Xi\|_{H^2} = O(\epsilon^2)$. ☐

5 Linear instability of small amplitude solitary waves

Our first observation here is that on spinor fields of the form

$$\psi(x,t)=\begin{bmatrix}\Psi_1(x,t)\\ \Psi_2(x,t)\end{bmatrix},\quad \psi(x,t)=\begin{bmatrix}\Psi_1(r,t)\\ ie^{i\phi}\Psi_2(r,t)\end{bmatrix},\quad \psi(x,t)=\begin{bmatrix}\Psi_1(r,t)\begin{pmatrix}1\\0\end{pmatrix}\\ i\Psi_2(r,t)\begin{pmatrix}\cos\theta\\ e^{i\phi}\sin\theta\end{pmatrix}\end{bmatrix} \tag{5.1}$$

in dimensions $n=1$, $n=2$, and $n=3$ respectively, the nonlinear Dirac equation (2.1) reduces to the system

$$\begin{cases} i\partial_t\Psi_1=(\partial_r+\frac{n-1}{r})\Psi_2+m\Psi_1-f(|\Psi_1|^2-|\Psi_2|^2)\Psi_1\ ,\\ i\partial_t\Psi_2=-\partial_r\Psi_1-m\Psi_2+f(|\Psi_1|^2-|\Psi_2|^2)\Psi_2\end{cases} \tag{5.2}$$

(with the convention $r=x$ for $n=1$). The solitary waves considered here lie in this class of fields, and so to prove their instability it suffices to show that they are unstable as solutions of (5.2).

5.1 Linearization at a solitary wave

To derive the linearization of system (5.2) at a solitary wave (2.4)-(2.5) we consider solutions in the form of the Ansatz

$$\begin{bmatrix}\Psi_1(r,t)\\ \Psi_2(r,t)\end{bmatrix}=\left(\begin{bmatrix}v(r,\omega)\\ u(r,\omega)\end{bmatrix}+\begin{bmatrix}\rho_1(r,t)\\ \rho_2(r,t)\end{bmatrix}\right)e^{-i\omega t}, \tag{5.3}$$

where

$$\rho(r,t):=\begin{bmatrix}\rho_1(r,t)\\ \rho_2(r,t)\end{bmatrix}\in\mathbb{C}^2.$$

Inserting this Ansatz into system (5.2) with $f(s)=s^k$ and recalling that $u(r,\omega)$ and $v(r,\omega)$ are real-valued, we find that the linearized system for ρ is

$$i\partial_t\rho=\begin{bmatrix}m-\omega-(v^2-u^2)^k & \partial_r+\frac{n-1}{r}\\ -\partial_r & -m-\omega+(v^2-u^2)^k\end{bmatrix}\rho-2k(v^2-u^2)^{k-1}\begin{bmatrix}v^2 & -uv\\ -uv & u^2\end{bmatrix}\operatorname{Re}\rho. \tag{5.4}$$

We note that the above equation is $\mathbb{R}$-linear but not $\mathbb{C}$-linear, due to the presence of $\operatorname{Re}\rho=\begin{bmatrix}\operatorname{Re}\rho_1\\ \operatorname{Re}\rho_2\end{bmatrix}$. We rewrite equation (5.4) in terms of $\operatorname{Re}\rho\in\mathbb{R}^2$ and $\operatorname{Im}\rho\in\mathbb{R}^2$:

$$\partial_t\begin{bmatrix}\operatorname{Re}\rho\\ \operatorname{Im}\rho\end{bmatrix}=\mathbf{J}\mathbf{L}(\omega)\begin{bmatrix}\operatorname{Re}\rho\\ \operatorname{Im}\rho\end{bmatrix} \tag{5.5}$$

where $\mathbf{J}$ corresponds to $1/i$:

$$\mathbf{J}=\begin{bmatrix}0 & I_2\\ -I_2 & 0\end{bmatrix},$$

and the 4×4 matrix operator $\mathbf{L}(\omega)$ is defined by

$$\mathbf{L}(\omega)=\begin{bmatrix}\mathsf{L}_+(\omega) & 0\\ 0 & \mathsf{L}_-(\omega)\end{bmatrix},$$

where, writing

$$f:=f(\phi_\omega^*\beta\phi_\omega)=(v^2(r,\omega)-u^2(r,\omega))^k,\quad f':=f'(\phi_\omega^*\beta\phi_\omega),$$

we have

$$\mathsf{L}_-(\omega)=\begin{bmatrix}m-\omega-f & \partial_r+\frac{n-1}{r}\\ -\partial_r & -m-\omega+f\end{bmatrix}=\begin{bmatrix}m-\omega-(v^2-u^2)^k & \partial_r+\frac{n-1}{r}\\ -\partial_r & -m-\omega+(v^2-u^2)^k\end{bmatrix}, \tag{5.6}$$

and

$$\begin{aligned}\mathsf{L}_+(\omega) &= \mathsf{L}_-(\omega) - 2f' \begin{bmatrix} v^2 & -uv \\ -uv & u^2 \end{bmatrix} \\ &= \begin{bmatrix} m-\omega-(v^2-u^2)^k - 2k(v^2-u^2)^{k-1}v^2 & \partial_r + \frac{n-1}{r} + 2k(v^2-u^2)^{k-1}uv \\ -\partial_r + 2k(v^2-u^2)^{k-1}uv & -m-\omega+(v^2-u^2)^k - 2k(v^2-u^2)^{k-1}u^2 \end{bmatrix}.\end{aligned} \tag{5.7}$$

Let us remind the reader that $v = v(r,\omega)$ and $u = u(r,\omega)$ in (5.6)-(5.7) both depend on ω.

Then equation (5.5) which describes the linearization of the reduced system (5.2) at the solitary wave $\phi_\omega e^{-i\omega t}$, takes the form

$$\partial_t \begin{bmatrix} \operatorname{Re}\rho \\ \operatorname{Im}\rho \end{bmatrix} = \mathbf{J}\mathbf{L}(\omega) \begin{bmatrix} \operatorname{Re}\rho \\ \operatorname{Im}\rho \end{bmatrix} = \begin{bmatrix} 0 & \mathsf{L}_-(\omega) \\ -\mathsf{L}_+(\omega) & 0 \end{bmatrix} \begin{bmatrix} \operatorname{Re}\rho \\ \operatorname{Im}\rho \end{bmatrix}.$$

For the sake of completeness we record here the essential spectrum of the linearized operator:

Lemma 5.1. $\sigma_{ess}(\mathbf{J}\mathbf{L}(\omega)) = i\mathbb{R}\backslash i(-1+|\omega|, 1-|\omega|)$.

Proof. The proof follows from noticing that, due to the exponential spatial decay of $v(r,\omega)$, $u(r,\omega)$ and the Weyl theorem, $\sigma_{ess}(\mathbf{J}\mathbf{L}(\omega)) = \sigma_{ess}(\mathbf{J}(\mathbf{D}_m - \omega))$, with $\mathbf{D}_m = \begin{bmatrix} D_m & 0 \\ 0 & D_m \end{bmatrix}$, $D_m = i\sigma_2\partial_r + \sigma_3 m$. At the same time, since $D_m^2 = -\Delta + m^2$, $\sigma_{ess}(D_m) = \mathbb{R}\backslash(-m,m)$, while $\mathbf{J}$ commutes with $\mathbf{D}_m$ and $\sigma(\mathbf{J}) = \{\pm i\}$, one concludes that

$$\sigma_{ess}(\mathbf{J}(\mathbf{D}_m - \omega)) = \sigma_{ess}(i(D_m-\omega)) \cup \sigma_{ess}(-i(D_m-\omega)) = i\mathbb{R}\backslash i(-m+|\omega|, m-|\omega|).$$

□

5.2 Unstable eigenvalue of $\mathbf{J}\mathbf{L}$ for $\omega \lesssim m$

Proposition 5.2. *Let $k \ge 3$ for $n = 1$, or $k \ge 2$ for $n = 2$, or $k = 1$ for $n = 3$. There is $\omega_0 < m$ (which depends on n and k) such that for $\omega \in (\omega_0, m)$ there are two families of eigenvalues*

$$\pm\lambda_\omega \in \sigma_p(\mathbf{J}\mathbf{L}(\omega)), \qquad \lambda_\omega > 0, \qquad \lambda_\omega = O(m-\omega).$$

Proof. The relation $\begin{bmatrix} 0 & \mathsf{L}_- \\ -\mathsf{L}_+ & 0 \end{bmatrix} \begin{bmatrix} \varphi \\ \vartheta \end{bmatrix} = \lambda \begin{bmatrix} \varphi \\ \vartheta \end{bmatrix}$, with φ, $\vartheta \in \mathbb{C}^2$, can be written explicitly as follows:

$$\begin{bmatrix} -\lambda & 0 & m-\omega-f & \partial_r + \frac{n-1}{r} \\ 0 & -\lambda & -\partial_r & -m-\omega+f \\ -m+\omega+f+2f'v^2 & -\partial_r - 2f'vu & -\lambda & 0 \\ \partial_r - 2f'vu & m+\omega-f+2f'u^2 & 0 & -\lambda \end{bmatrix} \begin{bmatrix} \varphi_1 \\ \varphi_2 \\ \vartheta_1 \\ \vartheta_2 \end{bmatrix} = 0. \tag{5.8}$$

We divide the first and the third rows by $\epsilon^2 = m^2 - \omega^2$, the second and the fourth rows by ϵ, and substitute $R = \epsilon r$, $\varphi_2 = \epsilon\Phi_2$, $\vartheta_2 = \epsilon\Theta_2$, to get

$$\begin{bmatrix} -\frac{\lambda}{\epsilon^2} & 0 & \frac{m-\omega-f}{\epsilon^2} & \partial_R + \frac{n-1}{R} \\ 0 & -\lambda & -\partial_R & -m-\omega \\ \frac{-m+\omega+f+2f'v^2}{\epsilon^2} & -(\partial_R + \frac{n-1}{R}) - \frac{2f'vu}{\epsilon} & -\frac{\lambda}{\epsilon^2} & 0 \\ \partial_R - \frac{1}{\epsilon}2f'vu & m+\omega-f+2f'u^2 & 0 & -\lambda \end{bmatrix} \begin{bmatrix} \varphi_1 \\ \Phi_2 \\ \vartheta_1 \\ \Theta_2 \end{bmatrix} = 0. \tag{5.9}$$

Anticipating the $\epsilon \to 0$ limit, formally set $\Lambda = \lim\limits_{\epsilon\to 0} \frac{\lambda}{\epsilon^2}$, and introduce the matrices

$$\mathbf{A}_\Lambda = \begin{bmatrix} -\Lambda & 0 & \frac{1}{2m} - \hat{V}^{2k}(R) & \partial_R + \frac{n-1}{R} \\ 0 & 0 & -\partial_R & -2m \\ -\frac{1}{2m} + (2k+1)\hat{V}^{2k}(R) & -(\partial_R + \frac{n-1}{R}) & -\Lambda & 0 \\ \partial_R & 2m & 0 & 0 \end{bmatrix}, \tag{5.10}$$

$$\mathbf{K}_1 = \mathrm{diag}[1,0,1,0], \qquad \mathbf{K}_2 = \mathrm{diag}[0,1,0,1], \tag{5.11}$$

where $\hat{V}(R)$, the NLS ground state, was introduced in (4.15). We write (5.9) in the form

$$\mathbf{A}_\Lambda \eta = \Big(\frac{\lambda}{\epsilon^2} - \Lambda\Big)\mathbf{K}_1\eta + \lambda\mathbf{K}_2\eta + W\eta, \qquad \eta = \begin{bmatrix} \varphi_1 \\ \varPhi_2 \\ \vartheta_1 \\ \varTheta_2 \end{bmatrix} \in \mathbb{C}^4, \tag{5.12}$$

where $W(R,\epsilon)$ is given by

$$\begin{bmatrix} 0 & 0 & \frac{1}{2m} - \hat{V}^{2k} + \frac{\omega-m+f}{\epsilon^2} & 0 \\ 0 & 0 & 0 & -m+\omega-f \\ -\frac{1}{2m} + (2k+1)\hat{V}^{2k} + \frac{m-\omega-f-2f'v^2}{\epsilon^2} & \frac{2f'vu}{\epsilon} & 0 & 0 \\ \frac{2f'vu}{\epsilon} & m-\omega+f+2f'u^2 & 0 & 0 \end{bmatrix} \tag{5.13}$$

is a zero order differential operator with L^∞ coefficients.

Lemma 5.3. $\|W(\cdot,\epsilon)\|_{L^\infty(\mathbb{R}^+,\mathbb{C}^4\to\mathbb{C}^4)} \le O(\epsilon^2)$.

Proof. By Lemma 4.2 (and the Sobolev inequality), one has

$$v(r,\omega) = \epsilon^{\frac{1}{k}}(\hat{V}(R) + \tilde{V}(R)), \qquad u(r,\omega) = \epsilon^{1+\frac{1}{k}}(\hat{U}(R) + \tilde{U}(R)), \qquad \|\tilde{V}\|_{L^\infty} + \|\tilde{U}\|_{L^\infty} = O(\epsilon^2).$$

Then

$$f(v^2-u^2) = \epsilon^2(V^2-U^2)^k,$$

$$f'(v^2-u^2) = k\epsilon^{2-\frac{2}{k}}(V^2-\epsilon^2U^2)^{k-1},$$

$$-\frac{1}{2m} + \frac{m-\omega-f-2f'v^2}{\epsilon^2} = O(\epsilon^2) - (V^2-\epsilon^2U^2)^k - 2kV^2(V^2-\epsilon^2U^2)^{k-1} = -(1+2k)\hat{V}^{2k} + O_{L^\infty}(\epsilon^2),$$

$$\frac{f'vu}{\epsilon} = k\epsilon^2UV(V^2-\epsilon U^2)^{k-1} = O_{L^\infty}(\epsilon^2),$$

$$m-\omega+f = O(\epsilon^2) + \epsilon^2(V^2-\epsilon^2U^2)^k = O_{L^\infty}(\epsilon^2),$$

$$f'u^2 = k\epsilon^4U^2(V^2-\epsilon U^2)^{k-1} = O(\epsilon^4)_{L^\infty},$$

and

$$\frac{1}{2m} + \frac{\omega-m+f}{\epsilon^2} = +O(\epsilon^2) + (V^2-\epsilon^2U^2)^k = \hat{V}^{2k} + O_{L^\infty}(\epsilon^2),$$

and the Lemma follows directly from this list of estimates. ☐

Lemma 5.4. $\dim\ker\mathbf{A}_\Lambda = \dim\ker(\mathbf{j}\mathfrak{l}_{rad} - \Lambda)$, *where*

$$\mathbf{j}\mathfrak{l}_{rad} = \begin{bmatrix} 0 & \mathfrak{l}_{+,rad} \\ -\mathfrak{l}_{-,rad} & 0 \end{bmatrix},$$

and where, we recall,

$$\mathfrak{l}_{-,rad} = -\frac{1}{2m}(\partial_R + \frac{n-1}{R})\partial_R + \frac{1}{2m} - \hat{V}(R)^{2k},$$

$$\mathfrak{l}_{+,rad} = -\frac{1}{2m}(\partial_R + \frac{n-1}{R})\partial_R + \frac{1}{2m} + (2k+1)\hat{V}(R)^{2k}.$$

Moreover, if $n=1$ and $k \ge 3$, or $n=2$ and $k \ge 2$, or if $n=3$ and $k=1$, there is $\Lambda > 0$ such that $\pm\Lambda \in \sigma_d(\mathbf{j}\mathbf{H})$ are simple eigenvalues.

Proof. An easy computation shows that

$$\Phi = \begin{bmatrix} \Phi_1 \\ \Phi_2 \\ \Phi_3 \\ \Phi_4 \end{bmatrix} \in \ker \mathbf{A}_\Lambda \quad \Longleftrightarrow \quad 2m\Phi_2 = -\partial_R \Phi_1, \quad 2m\Phi_4 = -\partial_R\Phi_3, \quad (\mathbf{j}\mathsf{l}_{rad} - \Lambda)\begin{bmatrix} \Phi_3 \\ -\Phi_1 \end{bmatrix} = 0,$$

and the first statement of the Lemma follows from this observation. The second statement then follows from Lemma 3.2. ☐

So we may assume that there is $\Lambda > 0$ such that $\pm\Lambda \in \sigma_d(\mathbf{A}_\Lambda)$, with eigenfunctions

$$\ker \mathbf{A}_{\pm\Lambda} \ni \Phi_{\pm\Lambda} = \begin{bmatrix} \pm\Phi_1 \\ \mp\frac{1}{2m}\partial_R\Phi_1 \\ \Phi_3 \\ -\frac{1}{2m}\partial_R\Phi_3 \end{bmatrix}, \quad \mathsf{l}_{+,rad}\Phi_1 = -\Lambda\Phi_3, \quad \mathsf{l}_{-,rad}\Phi_3 = \Lambda\Phi_1.$$

We will use the Rayleigh-Schrödinger perturbation theory to show that there are $\lambda \in \sigma_d(\mathbf{JL})$ with $\lambda \approx \pm\epsilon^2\Lambda$.

Writing

$$\mathbf{A}_\Lambda = \mathbf{J}\mathbf{L}_0 - \Lambda\mathbf{K}_1, \quad \mathbf{L}_0 = \begin{bmatrix} \mathsf{L}_{0,+} & 0 \\ 0 & \mathsf{L}_{0,-} \end{bmatrix},$$

with

$$\mathsf{L}_{0,+} = \begin{bmatrix} \frac{1}{2m} - (2k+1)\hat{V}^{2k} & \partial_R + \frac{n-1}{R} \\ -\partial_R & -2m \end{bmatrix}, \quad \mathsf{L}_{0,-} = \begin{bmatrix} \frac{1}{2m} - \hat{V}^{2k} & \partial_R + \frac{n-1}{R} \\ -\partial_R & -2m \end{bmatrix}$$

self-adjoint, we see that

$$\mathbf{A}_\Lambda^* = -\mathbf{L}_0\mathbf{J} - \Lambda\mathbf{K}_1 = \mathbf{F}\mathbf{A}_\Lambda\mathbf{F}, \quad \mathbf{F} := \begin{bmatrix} 0 & I_2 \\ I_2 & 0 \end{bmatrix}.$$

Hence $\ker \mathbf{A}_\Lambda^*$ is spanned by $\Phi_\Lambda^* := \mathbf{F}\Phi_\Lambda$. Let $\mathbf{P}_\Lambda$ denote the orthogonal projection onto Φ_Λ^*.

Seeking η and λ in the form

$$\eta = \Phi_\Lambda + \zeta, \quad \zeta \perp \Phi_\Lambda, \qquad \lambda = \epsilon^2(\Lambda + \mu),$$

then (5.12) becomes

$$\mathbf{A}_\Lambda \zeta = \mu\mathbf{K}_1(\Phi_\Lambda + \zeta) + \epsilon^2(\Lambda+\mu)\mathbf{K}_2(\Phi_\Lambda+\zeta) + W(\Phi_\Lambda+\zeta). \tag{5.14}$$

Applying $\mathbf{P}_\Lambda$ and $1 - \mathbf{P}_\Lambda$ to (5.14), one has:

$$0 = \mu\langle\Phi_\Lambda^*, \mathbf{K}_1(\Phi_\Lambda+\zeta)\rangle + \epsilon^2(\Lambda+\mu)\langle\Phi_\Lambda^*, \mathbf{K}_2(\Phi_\Lambda+\zeta)\rangle + \langle\Phi_\Lambda^*, W(\Phi+\zeta)\eta\rangle, \tag{5.15}$$

$$\mathbf{A}_\Lambda\zeta = (1-\mathbf{P}_\Lambda)\Big(\mu\mathbf{K}_1 + \epsilon^2(\Lambda+\mu)\mathbf{K}_2 + W\Big)(\Phi_\Lambda+\zeta). \tag{5.16}$$

Next, we claim

$$\langle\Phi_\Lambda^*, \mathbf{K}_1\Phi_\Lambda\rangle \neq 0. \tag{5.17}$$

To see this, note that

$$\langle\Phi_\Lambda^*, \mathbf{K}_1\Phi_\Lambda\rangle = \langle\mathbf{F}\Phi_\Lambda, \mathbf{K}_1\Phi_\Lambda\rangle = 2\,\mathrm{Re}\langle\Phi_3, \Phi_1\rangle.$$

Now the fact that $\Phi_\Lambda \in \ker \mathbf{A}_\Lambda$ means in particular that $L_-\Phi_3 = \Lambda\Phi_1$. Hence, since L_- is self-adjoint,

$$\mathbb{R} \ni \langle\Phi_3, L_-\Phi_3\rangle = \Lambda\langle\Phi_3, \Phi_1\rangle,$$

and so $\mathrm{Re}\langle\Phi_3,\Phi_1\rangle = 0$ only if $\langle\Phi_3, L_-\Phi_3\rangle = 0$. As is well-known, since $L_-\hat{V} = 0$ and $\hat{V}(r) > 0$, we have $L_- \geq 0$. Thus $\mathrm{Re}\langle\Phi_3,\Phi_1\rangle = 0$ only if $L_-\Phi_3 = 0$. This, in turn, would imply either $\Lambda = 0$ or $\Phi_\Lambda = 0$, both of which are false.

Denote by $L^2_r(\mathbb{R}^n,\mathbb{C}^4) \subset L^2(\mathbb{R}^n,\mathbb{C}^4)$ the subspace of spherically symmetric functions. Now using (5.17), and the existence of the bounded inverse $\mathbf{A}_\Lambda^{-1} : \mathrm{Range}\,(1-\mathbf{P}_\Lambda) \to \Phi_\Lambda^\perp$, the equations (5.15), (5.16) can be written as

$$\mu = M(\mu,\zeta), \qquad \zeta = Z(\mu,\zeta),$$

with functions $M : \mathbb{R}\times L^2_r \to \mathbb{R}$, $\ Z : \mathbb{R}\times L^2_r \to L^2_r$ given by

$$M(\mu,\zeta) = -\frac{1}{\langle \Phi_\Lambda^*, \mathbf{K}_1\Phi_\Lambda\rangle}\Big[\mu\langle\Phi_\Lambda^*,\mathbf{K}_1\zeta\rangle + \langle\Phi_\Lambda^*, \epsilon^2(\Lambda+\mu)\mathbf{K}_2(\Phi_\Lambda+\zeta)+W\rangle\Big], \tag{5.18}$$

$$Z(\mu,\zeta) = \mathbf{A}_\Lambda^{-1}(1-\mathbf{P}_\Lambda)\Big(\mu\mathbf{K}_1 + \epsilon^2(\Lambda+\mu)\mathbf{K}_2 + W\Big)(\Phi_\Lambda+\zeta). \tag{5.19}$$

Pick $\Gamma \geq 1$ such that

$$\Gamma \geq 2\|\mathbf{A}_\Lambda^{-1}(1-\mathbf{P}_\Lambda)\mathbf{K}_1\Phi_\Lambda\|_{L^2}. \tag{5.20}$$

Lemma 5.5. *Consider* $\mathbb{R}\times L^2_r$ *endowed with the metric*

$$\|(\mu,\zeta)\|_\Gamma = \Gamma|\mu| + \|\zeta\|_{L^2}.$$

There is $\omega_1 \in (\omega_0, m)$ *such that for* $\omega \in (\omega_1, m)$ *the map*

$$M\times Z :\ \ \mathbb{R}\times L^2_r \to \mathbb{R}\times L^2_r, \qquad (\mu,\zeta)\mapsto \big(M(\mu,\zeta), Z(\mu,\zeta)\big), \tag{5.21}$$

restricted onto the set

$$\mathcal{B}_\epsilon = \{(\mu,\zeta)\in\mathbb{R}\times L^2(\mathbb{R},\mathbb{C}^4);\ \|(\mu,\zeta)\|_\Gamma \leq \epsilon\} \subset \mathbb{R}\times L^2_r$$

is an endomorphism and a contraction with respect to $\|\cdot\|_\Gamma$.

Proof. Assuming $\Gamma|\mu| + \|\zeta\|_{L^2} \leq \epsilon < 1$, and using Lemma 5.3, we have the estimates

$$|M(\mu,\zeta)| \leq C\left(|\mu|\|\zeta\|_{L^2} + \epsilon^2(1+|\mu|)(1+\|\zeta\|_{L^2}) + \|W\|_{L^\infty}\right) \leq C\epsilon^2,$$

$$\|Z(\mu,\zeta)\|_{L^2} \leq \frac{1}{2}\Gamma|\mu| + C\left(|\mu|\|\zeta\|_{L^2} + \epsilon^2(1+|\mu|)(1+\|\zeta\|_{L^2}) + \|W\|_{L^\infty}(1+\|\zeta\|_{L^2})\right) \leq \frac{1}{2}\epsilon + C\epsilon^2$$

which show that for all sufficiently small ϵ, $M\times Z$ maps $\mathcal{B}_\epsilon$ into itself. Similar estimates show that $(M\times Z)|_{\mathcal{B}_\epsilon}$ is a contraction in the metric $\|\cdot\|_\Gamma$. ☐

According to Lemma 5.5, by the contraction mapping theorem, for $\omega \in (\omega_1, m)$, the map (5.21) has a unique fixed point $(\mu_0(\omega),\zeta_0(\omega)) \in \mathcal{B}_\epsilon \subset \mathbb{R}\times L^2_r$. Thus, we have $\pm\epsilon^2(\Lambda+\mu_0(\omega)) \in \sigma_p(\mathbf{JL}(\omega))$, $\omega\in(\omega_1,m)$, with $\Gamma|\mu_0(\omega)| \leq \epsilon$, finishing the proof of the proposition. ☐

By Remark 2.3, Proposition 5.2 finishes the proof of Theorem 2.1.

References

[AS83] A. Alvarez and M. Soler, Phys. Rev. Letters, 50 (1983), pp. 1230–1233.

[AS86] A. Alvarez and M. Soler, Phys. Rev. D, 34 (1986), pp. 644–645.

[BC09] G. Berkolaiko and A. Comech, *On spectral stability of solitary waves of nonlinear Dirac equation on a line*, ArXiv e-prints, 0910.0917 (2009).

[BC11] N. Boussaid and S. Cuccagna, *On stability of standing waves of nonlinear Dirac equations*, ArXiv e-prints, 1103.4452 (2011).

[BL83] H. Berestycki and P.-L. Lions, *Nonlinear scalar field equations. I. Existence of a ground state*, Arch. Rational Mech. Anal., 82 (1983), pp. 313–345.

[Bou08] N. Boussaid, *On the asymptotic stability of small nonlinear Dirac standing waves in a resonant case*, SIAM J. Math. Anal., 40 (2008), pp. 1621–1670.

[Chu07] M. Chugunova, *Spectral stability of nonlinear waves in dynamical systems (Doctoral Thesis)*, McMaster University, Hamilton, Ontario, Canada, 2007.

[CKMS10] F. Cooper, A. Khare, B. Mihaila, and A. Saxena, *Solitary waves in the nonlinear Dirac equation with arbitrary nonlinearity*, Phys. Rev. E, 82 (2010), p. 036604.

[Com11] A. Comech, *On the meaning of the Vakhitov-Kolokolov stability criterion for the nonlinear Dirac equation*, ArXiv e-prints, 1107.1763 (2011).

[CP03] A. Comech and D. Pelinovsky, *Purely nonlinear instability of standing waves with minimal energy*, Comm. Pure Appl. Math., 56 (2003), pp. 1565–1607.

[CP06] M. Chugunova and D. Pelinovsky, *Block-diagonalization of the symmetric first-order coupled-mode system*, SIAM J. Appl. Dyn. Syst., 5 (2006), pp. 66–83.

[CV86] T. Cazenave and L. Vázquez, *Existence of localized solutions for a classical nonlinear Dirac field*, Comm. Math. Phys., 105 (1986), pp. 35–47.

[Dir28] P. A. M. Dirac, *The quantum theory of the electron*, Proc. Roy. Soc. A., 117 (1928), pp. 616–624.

[ES95] M. J. Esteban and É. Séré, *Stationary states of the nonlinear Dirac equation: a variational approach*, Comm. Math. Phys., 171 (1995), pp. 323–350.

[GN74] D. J. Gross and A. Neveu, *Dynamical symmetry breaking in asymptotically free field theories*, Phys. Rev. D, 10 (1974), pp. 3235–3253.

[Gro66] L. Gross, *The Cauchy problem for the coupled Maxwell and Dirac equations*, Comm. Pure Appl. Math., 19 (1966), pp. 1–15.

[GSS87] M. Grillakis, J. Shatah, and W. Strauss, *Stability theory of solitary waves in the presence of symmetry. I*, J. Funct. Anal., 74 (1987), pp. 160–197.

[Gua08] M. Guan, *Solitary wave solutions for the nonlinear Dirac equations*, ArXiv e-prints, 0812.2273 (2008).

[HC09] L. H. Haddad and L. D. Carr, *The nonlinear Dirac equation in Bose-Einstein condensates: foundation and symmetries*, Phys. D, 238 (2009), pp. 1413–1421.

[LG75] S. Y. Lee and A. Gavrielides, *Quantization of the localized solutions in two-dimensional field theories of massive fermions*, Phys. Rev. D, 12 (1975), pp. 3880–3886.

[Mer88] F. Merle, *Existence of stationary states for nonlinear Dirac equations*, J. Differential Equations, 74 (1988), pp. 50–68.

[MJZ$^+$10] M. Merkl, A. Jacob, F. E. Zimmer, P. Öhberg, and L. Santos, *Chiral confinement in quasirelativistic Bose-Einstein condensates*, Phys. Rev. Lett., 104 (2010), p. 073603.

[Pau36] W. Pauli, *Contributions mathématiques à la théorie des matrices de Dirac*, Ann. Inst. H. Poincaré, 6 (1936), pp. 109–136.

[Pel10] D. Pelinovsky, *Survey on global existence in the nonlinear Dirac equations in one dimension*, ArXiv e-prints, 1011.5925 (2010).

[PS10] D. E. Pelinovsky and A. Stefanov, *Asymptotic stability of small gap solitons in the nonlinear Dirac equations*, ArXiv e-prints, 1008.4514 (2010).

[Sha83] J. Shatah, *Stable standing waves of nonlinear Klein-Gordon equations*, Comm. Math. Phys., 91 (1983), pp. 313–327.

[Sol70] M. Soler, *Classical, stable, nonlinear spinor field with positive rest energy*, Phys. Rev. D, 1 (1970), pp. 2766–2769.

[SS85] J. Shatah and W. Strauss, *Instability of nonlinear bound states*, Comm. Math. Phys., 100 (1985), pp. 173–190.

[Str77] W. A. Strauss, *Existence of solitary waves in higher dimensions*, Comm. Math. Phys., 55 (1977), pp. 149–162.

[SV86] W. A. Strauss and L. Vázquez, *Stability under dilations of nonlinear spinor fields*, Phys. Rev. D (3), 34 (1986), pp. 641–643.

[Tha92] B. Thaller, *The Dirac equation*, Texts and Monographs in Physics, Springer-Verlag, Berlin, 1992.

[Thi58] W. E. Thirring, *A soluble relativistic field theory*, Ann. Physics, 3 (1958), pp. 91–112.

[vdW32] B. L. van der Waerden, *Die gruppentheoretische Methode in der Quantenmechanik*, Springer-Verlag, Berlin, 1932.

[VK73] N. G. Vakhitov and A. A. Kolokolov, *Stationary solutions of the wave equation in the medium with nonlinearity saturation*, Radiophys. Quantum Electron., 16 (1973), pp. 783–789.

[Wei85] M. I. Weinstein, *Modulational stability of ground states of nonlinear Schrödinger equations*, SIAM J. Math. Anal., 16 (1985), pp. 472–491.